


# FIPRONIL 0.05 % GEL



ROACH PROBLEM?

WE HAVE GOT SOLUTION.

**SPECIAL FORMULATION FOR ELIMINATING NASTY COCKROACHES**

**Syringe 35gm**

Fipronil 0.05 % gel is a household insecticide recommended for the control of American Cockroach (*Periplaneta Americana*) and German cockroach (*Blattella germanica*). Extremely low amount of the chemical is required to achieve the control. The application of Fipronil 0.05% w/w Gel is fast and convenient, as it causes no disturbance or discomfort and there is no need to leave premise during or after the treatments. It is highly effective, the most modern technology for cockroach control.

## RECOMMENDATIONS:

Fipronil 0.05 % gel is a household insecticide recommended for the control of American Cockroach (*Periplaneta americana*) and German cockroach (*Blattella germanica*). The bait gun dispenses 0.03g of Fipronil Gel at every shot 3 to 4 spots per square meter is recommended for good control.

## DIRECTION FOR USE:

1. Take the lid off from the end of the syringe bottle.
2. Place about 0.5g of the product (on paper or Aluminium foil) at 35 cm interval on harborage, corridor, wall voids, corners or edges of the floor, cabinet shelves etc where cockroaches may be a problem.
3. For more effective control of cockroaches use up all contents at a time and as its effect lasts 3 months. It is recommended to re-install after 2-3 months for continuous cockroaches control.
4. It is easy to install and gives excellent attractancy from special bait ingredients, so it controls small and large cockroaches effectively.

## CAUTIONS ON USE:

1. Read and follow all the instructions on the label and leaflet very carefully before using.
2. Avoid direct contact or inhalation of product by wearing face mask, rubber gloves, goggles, protective clothing and gum boots.
3. Do not eat, drink, chew or smoke while handling.
4. Wash contaminated clothes and parts of the body after application. Wear full protective clothing during application.
5. Call a physician immediately if any symptoms of poisoning appear in a person who has been handling the material.

## SYMPTOMS OF POISONING:

Vomiting, agitation, seizures, dizziness and paresthesia.

## FIRST AID MEASURES:

**If ingested:** If Person is conscious and alert, rinse mouth with water and give 1 or 2 glass of water to drink. Never give anything by mouth to an unconscious persons. Call the doctor. Don't induced vomiting until and unless desired by attending physician.

**If inhaled:** Take the patient to fresh air and take care of body temperature. If patient is not breathing or breathing with difficulty, give artificial respiration, preferably with mouth to mouth breathing. Consult a physician.

**In case of skin contact:** Wash thoroughly the affected part of body with soap and plenty of water including hairs and under fingernails. If pain, redness or irritation persists, consult a doctor. All the contaminated clothes should be segregated and thoroughly washed with soap and water separately before use.

**In case of eye contact:** Rinse gently with plenty of water at least for 15 minutes. If pain, irritation, redness or photophobia persists. Consult eye specialist.

**ANTIDOTE:**

No specific antidote. Treat symptomatically.

**STORAGE CONDITION:**

Air tight bottle, cool place

**CONTENT (ACTIVE INGREDIENT):**

FIPRONIL ..... 0.05% w/w



**KEEP OUT OF REACH OF CHILDREN AND PETS DURING INSTALLATION AND STORAGE AND AS IT CAN BE DANGEROUS WHEN CHILDREN PLAY WITH DISPOSE THE USED CONTAINER SEPARATELY**

**MANUFACTURED BY:**

Kukbo Science Co., Ltd.  
49, Sandon-RO, Hungdeok-KU, Cheongju, Chungbuk, Korea

**IMPORTED & MARKETING BY:**

Mahamaya Lifesciences Pvt. Ltd.  
Unit No: DPT – 033, DLF Prime Tower, F – Block, Okhla, Phase – 1,  
New Delhi – 110020, India  
[www.mahamayalifesciences.com](http://www.mahamayalifesciences.com)